

POST TEST

Improving Recognition and Prevention of Residual Paralysis After Surgery: Case Examples of Monitoring and Reversal

Name: Click here to enter text.		Credential(s): Click here to enter text.	
Address: Click here to enter text.			
City: Click here to enter text.	State: Click here to enter text.	Zip Code: Click here to enter text.	
Home Phone: Click here to enter text.		Work/Cell Phone: Click here to enter text.	
Email Address: Click here to enter text.			
AAMA Member: <input type="checkbox"/> Yes <input type="checkbox"/> No Member# Click here to enter text.		Visa Mastercard AMEX Discover Card# _____ EXP: _____ Code: _____	

Deadline September 30, 2017

Please email test to ceuprograms@ctsma.org

Or mail to Jennifer Amaral CMA(AAMA) 208 Deerfield Ridge Dr, Mystic CT 06355

Make Checks Payable to CSMA.

POST TEST

Improving Recognition and Prevention of Residual Paralysis After Surgery: Case Examples of Monitoring and Reversal

1. The incidence of RNMB is much higher than anesthesiologists report in surveys.
 - True
 - False
2. A meta-analysis of 24 studies that included 3,375 patients found that ____% of patients had a TOFR less than 0.9.
 - 10
 - 20
 - 40
 - 80
3. RNMB is associated with which of the following postoperative complications?
 - Critical respiratory events
 - Longer duration of hospital stay
 - Poorer quality of recovery
 - All of the above
4. Which of the following is the most reliable method for monitoring RNMB?
 - Visual assessment of the adductor pollicis
 - Tactile assessment of the orbicularis oculi
 - Objective measurement of the adductor pollicis
 - Clinical tests of muscle weakness such as head lift or sustained tongue depressor test
5. Which of the following statements about sugammadex dosing is true?
 - 2 mg/kg is appropriate if recovery has reached the reappearance of the second twitch in response to TOF stimulation.
 - 4 mg/kg is appropriate if recovery has reached 1 to 2 post-tetanic counts and no twitch response to TOF stimulation is observed.
 - Doses should be calculated according to actual or total body weight in obese patients.
 - All of the above
6. How long after the administration of neostigmine should tracheal extubation occur?
 - 10 minutes
 - 15 minutes
 - 20 minutes
 - None of the above

7. Which of the following statements about ACh in the NMJ is true?

- ACh must bind both α subunits on the AChR for depolarization of the muscle fiber to occur.
- All AChRs on the muscle fiber endplate require binding to generate an action potential.
- Precisely the amount of ACh needed to bind the AChRs is released from the motor neuron into the synaptic cleft.
- All of the above

8. Which of the following statements about NMBAs is true?

- NMBAs must bind both α subunits to prevent depolarization of the muscle fiber.
- More than 90% of the AChRs must be occupied by the NMBA to completely block neuromuscular transmission.
- All of the above
- None of the above